



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Colymbus auritus. HORNE GREBE.—Very abundant during the migrations during last fall and this spring. I first observed them October 18, 1902, near Fair Haven, on Lake St. Clair. In April, of this year, I found them common in the Detroit River above the city. On the 27th I saw about fifty birds, on May 4 about sixty. They were generally unsuspicious and allowed a near approach. I saw the last May 10, twelve birds.—BRADSHAW H. SWALES, *Detroit, Mich.*

Occurrence of the Ruff (*Pavoncella pugnax*) and Other Birds in Rhode Island. — **Larus atricilla.** LAUGHING GULL.—I observed two birds of this species on a marsh at Seaconnet Point on Aug. 24, 1903. One of the birds was in adult plumage, but the other seemed immature. This species is not often seen in Rhode Island, there being but one instance of its capture in the State recorded in 'The Birds of Rhode Island' by Howe and Sturtevant.

Hydrochelidon nigra surinamensis. BLACK TERN.—A fine male of this species was shot near Newport on July 30, 1903. It was just beginning to lose the black plumage.

Ardetta exilis. LEAST BITTERN.—A bird of this species was shot on July 18, 1903, on a salt marsh near Newport. It is now in my collection. The Least Bittern, although formerly common near Newport, seems to have become rare during the last five years.

Micropalama himantopus. STILT SANDPIPER.—This species occurred in greater numbers than usual near Newport in August and early September, 1903. It seems to be a very irregular migrant, varying in numbers from year to year.

Limosa hæmastica. HUDSONIAN GODWIT.—Eighteen 'Ring-tail Marlins' were observed at Point Judith on August 30, 1903, and six were shot. Three of these latter, which I obtained, proved to be adult birds, two being males and one a female. They were changing into winter plumage but still had many traces of the reddish summer plumage on the breasts and flanks. The birds were seen during a severe northeast gale and were easily approached as they stood huddled together in a pool of water about five inches deep. This species is rare in Rhode Island, not more than one or two being shot each year.

Pavoncella pugnax. RUFF.—An immature female of this species was taken at Point Judith, R. I., on August 31, 1903, by a local gunner. I obtained it of him and it is now in my collection. The bird, which was flying alone, was shot over decoys. I believe this is the second record for this bird in Rhode Island.—LeROY KING, *Newport, R. I.*

The Black-bellied Plover, Road-runner, and Black-throated Green Warbler in Kansas.—I wish to restore to my 'Catalogue of the Birds of Kansas' the Black-bellied Plover (*Charadrius squatarola*). It was omitted from my 5th edition (May, 1903) because I had no personal